


HOPA Annual Conference 2025

Research Abstracts



HOPA

Together we can reach new heights



Completed Research Abstracts

HOPA Annual Conference 2025

Together we can reach new heights

Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

Characteristics and Dosing Patterns of US Patients Diagnosed With HR-Positive/HER2-Negative Early Breast Cancer Initiating Abemaciclib at a Lower Dose Than the Approved 150-mg Twice Daily

Presenting Author: Andrea LeVoir, PharmD Eli Lilly and Company, Indianapolis, IN

Co-Authors: Laura Spring, MD, Massachusetts General Hospital, Boston, MA; Wambui Gathirua-Mwangi, PhD, Sangmi Kim, PhD Zhanglin Lin Cui, PhD Samuel Whipple, MPH Astra M Liepa, PharmD Erich Brechtelsbauer, PharmD, MS Raisa Volodarsky, PharmD and Sarah Rybowski, MBBSEli Lilly and Company, Indianapolis, IN; Madeline Richey, MPH and Jingru Wang, BS Flatiron Health, New York City, NY; Hatem Soliman, MD Moffitt Cancer Center and Research Institute, Tampa, FL

BACKGROUND: Abemaciclib in combination with endocrine therapy is approved for adjuvant treatment of adult patients with hormone receptor–positive/HER2-negative, node-positive, early breast cancer at high risk of recurrence. Anecdotal insights indicate providers may initiate abemaciclib at a lower dose than the approved 150 mg twice daily to aid tolerability with the aim to increase to the approved dose. This real-world study describes demographic and clinical characteristics, dosing patterns, and incidence of prespecified adverse events (AEs) in patients with early breast cancer who were started on abemaciclib at a lower dose.

METHODS: This study used the nationwide Flatiron Health electronic health records–derived deidentified database. Patients started on abemaciclib October 2021 to November 2022 were included. All results were summarized descriptively. Additional analyses were conducted to compare characteristics of patients starting <150 mg twice daily versus 150 mg twice daily.

RESULTS: Among 453 patients, 65 started abemaciclib at <150 mg twice daily (median follow-up time 7.1 months). Median age was 56 years, 33.8% had stage III disease, 49.2% had ECOG Performance Scale score of 0 and median Charlson Comorbidity Index score was 0. Abemaciclib was frequently combined with aromatase inhibitors (92.3%), and most patients (56.9%) had a dose modification. The most common starting doses were 100 mg twice daily (56.9%) and 150 mg daily (24.6%); the time to first dose increase was 71 and 15 days, respectively. At first dose

modification, 32.3% patients increased dose to 150 mg twice daily. Common real world AEs were diarrhea (58.5%), fatigue (52.3%), and nausea/vomiting (40.0%). In comparison to patients started on 150 mg twice daily (n=388), patients started on a lower dose were more likely to be aged ≥ 75 years (4.9% vs 13.8%; $P=.022$).

CONCLUSIONS: In the real world, patients with hormone receptor–positive/HER2-negative early breast cancer started on abemaciclib at <150 mg twice daily were started on different doses and titration intervals. One-third of patients received dose increase to 150 mg twice daily. Real world AEs were generally consistent with clinical trial results.

This abstract was previously presented at the 2024 European Society for Medical Oncology Breast Cancer Annual Meeting.

HOPAC

Together we can reach new heights

Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

Clinical Characteristics and Treatment Persistence in US Patients With HR-Positive/HER2-Negative, Node Positive Early Breast Cancer Treated With Abemaciclib: Real-World Study From First Year After Approval

Presenting Author: Erich Brechtelsbauer, PharmD, MS, Eli Lilly and Company, Indianapolis, IN

Co-Authors: Kathryn Hudson MD, Texas Oncology, Austin, TX; Wambui Gathirua-Mwangi, PhD, Zhanglin Lin Cui, Brenda Grimes, PhD, Astra M Liepa, PharmD, Raisa Volodarsky, PharmD, and Katheryn Moreira, MD, MBA, Eli Lilly and Company, Indianapolis, IN; Madeline Richey, MPH and Jingru Wang, Flatiron Health, New York City, NY; Hatem Soliman, MD, Moffitt Cancer Center and Research Institute, Tampa, FL

BACKGROUND: Abemaciclib in combination with endocrine therapy is approved for adjuvant treatment of adult patients with hormone receptor–positive/HER2-negative, node-positive, early breast cancer at high risk of recurrence. The monarchE trial established the efficacy of abemaciclib and endocrine therapy for early breast cancer, with the highest rates of early discontinuation observed in the first few months. The use of abemaciclib in the real-world early breast cancer setting can provide insight into treatment patterns and inform adverse event (AE) management strategies. This retrospective study describes clinical characteristics and treatment persistence in patients with hormone receptor–positive/HER2-negative, node-positive early breast cancer starting abemaciclib.

METHODS: Data were accessed from Flatiron Health electronic database. Adult patients with node-positive, stage I-III early breast cancer starting abemaciclib October 2021 (FDA approval) to November 2022 at 150 mg twice daily were analyzed. Persistence rate was defined as the percentage of patients remaining on abemaciclib at 3 months, allowing for ≤60-day medication gap.

RESULTS: A cohort of 354 patients with a median follow-up time from abemaciclib initiation of 8.8 months were selected. The median age was 56 years, 25.4% were aged ≥65 years, 12.7% were Black, 4.0% were Asian, and most patients (80.8%) received care in a community setting.

More than half (55.4%) patients were postmenopausal; 57.9% had an ECOG Performance Status 0; whereas 25.1% had ECOG Performance Status 1. Approximately 33.9% had ≥ 1 comorbidity, and 12.1% had ≥ 2 comorbidities, with diabetes (14.1%) being the most frequent. Most patients had stage II (41.8%) or III (38.4%) disease, nodal status N1 (45.2%) or N2 (35.3%), and tumor grade 2 (52.3%). Abemaciclib was initiated at a median of 11.1 months after early breast cancer diagnosis. Before abemaciclib initiation, most patients received radiotherapy (96.3%) and chemotherapy (83.1%), with 46.3% receiving neoadjuvant chemotherapy. Most patients (74.0%) initiated endocrine therapy 1.6 months before abemaciclib initiation. The median time to abemaciclib initiation from breast surgery was 6.7 months. The most frequent regimen was abemaciclib and aromatase inhibitors (91.0%). At 3 months, 81.6% of patients were persistent; 5.6% resumed abemaciclib after >60-day interruption and 11.3% discontinued due to AEs. Additional information on dose modifications will be presented.

CONCLUSION: In this real-world study of utilization of abemaciclib in the first year after approval for early breast cancer, an older, less fit, and more racially diverse population than participated in the monarchE trial, as well as a higher proportion of patients with lower nodal status, was observed. The high 3-month persistence rate suggests abemaciclib for early breast cancer is well-tolerated in routine clinical practice.

This abstract was previously presented at the 2024 San Antonio Breast Cancer Symposium

Together we can reach new heights

Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

Clinical Outcomes of 2-Dose Tandem Influenza Vaccination Strategy in Patients With Plasma Cell Dyscrasias

Presenting Authors: E. Bridget Kim, PharmD, BCPS, BCOP, Jack Malespini, PharmD, BCOP, Victoria Yeung, Brenna Rowen, PharmD, BCOP, and Jennifer A. Hutchinson, PharmD, BCOP, Massachusetts General Hospital, Boston, MA

BACKGROUND: Patients with multiple myeloma and other plasma cell dyscrasias (PCD) are especially susceptible to infections, including influenza (flu). Multiple risk factors arise from disease-related immunoparesis, alteration in T-cell immunity, treatment-related immunosuppression, diminished protection from flu vaccines, and age-related decline in immune function. After the standard flu vaccine, these patients have shown to have low hemagglutination antibody inhibition titers, with decreased seroprotection rates of <20%. A booster dose could enhance immune efficacy. At Massachusetts General Hospital, we began implementing a 2-dose tandem flu vaccination strategy in patients with PCD in January 2021. Although it was previously reported that tandem flu vaccines can improve seroprotection rates, it has not yet been linked to better clinical outcomes.

OBJECTIVE: To compare flu-related clinical outcomes in patients with PCD who received 1 versus 2 flu vaccine doses.

METHODS: This study was a retrospective review of patients with PCD who received ≥ 1 flu vaccine over the past 4 flu seasons in 2020-2024. The primary outcomes were documented flu infections, level of clinical care required, and duration of hospitalization/PCD treatment interruptions. Secondary outcomes included flu treatment and time from most recent vaccine to flu infection.

RESULTS: A total of 100 randomly selected patients with PCD were included. The majority of patients had multiple myeloma (80%), and other PCDs included monoclonal gammopathy of undetermined significance, AL amyloidosis, and Waldenstrom macroglobulinemia. We identified 58 cases of documented flu infections (6 confirmed; 52 possible). In comparing flu rates, 84% of

the flu cases occurred in patients who received 1 dose versus 16% in patients who received 2 doses of flu vaccine. The level of care required included phone call to the office (57% vs 5%), office visit (19% vs 7%), urgent care visit (7% vs 2%), hospitalization (2%, 30-day vs 2%, 10-day) in 1-dose versus 2-dose cohorts, respectively. PCD treatment interruption was required in 21% in 1-dose cohort versus 2% in 2-dose cohort. The mean duration of treatment interruption was 12 days (range, 1-30) and 10 days (range, none; **1 instance**) in 1-dose versus 2-dose cohorts, respectively. Oseltamivir was prescribed in 3.4% in 1-dose cohort versus 1.7% in 2-dose cohort. The time of flu from most recent vaccine was longer in the 1-dose cohort, with 237 days (range, 34-583), compared with 141 days (range, 52-420) in the 2-dose cohort.

CONCLUSIONS: The 2-dose tandem flu vaccines improve seroprotection and directly result in favorable flu-related clinical outcomes in patients with PCD. Other benefits include lower utilization of healthcare resources, decreased hospital stay, and decreased incidents of PCD treatment interruptions.

1. Branagan AR, Duffy E, Albrecht RA, et al. Clinical and serologic responses after a two-dose series of high-dose influenza vaccine in plasma cell disorders: a prospective, single-arm trial. *Clin Lymphoma Myeloma Leuk*. 2017;17:296-304.
2. Branagan A, Duffy E, Foster C, et al. Two dose series of high-dose influenza vaccine is associated with longer duration of serologic immunity in patients with plasma cell disorders. *Blood*. 2017;130(suppl 1):438.

Together we can reach new heights

Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

Dexamethasone Versus Tocilizumab for Management of Cytokine Release Syndrome Related to Teclistamab in Patients With Relapsed/Refractory Multiple Myeloma

Presenting Authors: James A. Davis, PharmD, BCOP, Katelynn Granger, PharmD, and Kimberly M. Green, DO, Hollings Cancer Center, The Medical University of South Carolina, Charleston, SC; Jordan Snyder, PharmD, BCOP, and Zahra Mahmoudjafari, PharmD, MBA, BCOP, FHOPA, University of Kansas Cancer Center, Westwood, KS; Mikhaila Rice, PharmD, BCOP, Jack Khouri, MD, and Joslyn Rudoni, PharmD, BCOP, Taussig Cancer Center, Cleveland Clinic, Cleveland, OH; Donald C Moore, PharmD, BCOP, Shebli Atrash, MD, Hailey Hill, PharmD, BCOP, Jessica McElwee, PharmD, BCOP, and Grace Elsey, PharmD, BCOP, Levine Cancer Institute, Charlotte, NC; Christopher Cahoon, PharmD, Kelley Julian, PharmD, BCOP, and Charlotte B. Wagner, PharmD, BCOP, Huntsman Cancer Institute, University of Utah, Salt Lake City, Utah; Victoria Nachar, PharmD, BCOP, University of Michigan Rogel Cancer Center, Ann Arbor, MI

BACKGROUND: Teclistamab is a B-cell maturation antigen-directed bispecific antibody approved for use in patients with relapsed/refractory multiple myeloma. In the MajesTEC-1 trial, cytokine release syndrome (CRS) occurred in 72.1% of patients and was primarily managed with tocilizumab.¹ Recently, the International Myeloma Working Group released guidelines supporting the use of tocilizumab for low-grade CRS.²

OBJECTIVE: To evaluate the safety and efficacy of CRS management with dexamethasone compared with tocilizumab.

METHODS: In all, 6 US academic medical centers contributed data on 243 patients who started teclistamab as of July 2024. All patients received teclistamab in a step-up dosing manner per the prescribing information. Patients received adverse event management based on institutional protocols. CRS was graded using the American Society for Transplantation and Cellular Therapy criteria.³ Outcomes included incidence and severity of CRS, overall response rate (ORR), progression-free survival (PFS), and overall survival (OS).

RESULTS: Of the 243 patients included, 133 (55%) patients had CRS (41% grade 1; 13% grade 2). Three grade 3+ events resolved with intensive care. At a median follow-up of 8.7 months, the ORR, median PFS, and median OS were 66%, 6, and 14.3 months, respectively. Dexamethasone was used 23% of patients, tocilizumab in 29%, and dexamethasone and tocilizumab were used in 23% of patients. Of the 30 patients in the dexamethasone and tocilizumab group, 11 received dexamethasone before tocilizumab. Of those 11, 6 received dexamethasone for immune effector cell-associated neurotoxicity syndrome and 2 of the remaining 5 received subsequent tocilizumab due to grade 2 CRS not resolved with dexamethasone.

Of the 31 patients who had CRS in the dexamethasone group, 87% had grade 1 events. Thirteen patients (42%) who received dexamethasone had recurrent CRS after a subsequent teclistamab dose; however, all subsequent events were of the same or lower grade and resolved with a median of 2 (range, 2-5) additional dexamethasone doses.

In the tocilizumab group, 68% and 32% of patients had grade 1 and 2 events, respectively. The majority (92%) of patients received 1 dose of tocilizumab, with 3 patients requiring multiple doses. Six patients (16%) had recurrent CRS after a subsequent dose.

Median duration of CRS was 1 day for both groups, and step-up doses were delayed a median of 1 day in approximately a quarter of patients in both groups ($P=.78$). ORR, PFS, and OS were similar between the groups.

CONCLUSIONS: Although more patients had subsequent CRS with dexamethasone, CRS was low grade and manageable with repeated dexamethasone. While considering advantages over tocilizumab in regard to availability, ease of use, and cost, this study highlights the feasibility of dexamethasone for the management of CRS in patients receiving teclistamab.

1. Moreau P, Garfall AL, van de Donk NWCJ, et al. Teclistamab in relapsed or refractory multiple myeloma. *N Engl J Med*. 2022;387:495-505.
2. Rodriguez-Otero P, Usmani S, Cohen AD, et al. International Myeloma Working Group immunotherapy committee consensus guidelines and recommendations for optimal use of T-cell-

engaging bispecific antibodies in multiple myeloma. *Lancet Oncol.* 2024;25:e205-e216. Erratum in: *Lancet Oncol.* 2024;25:e284.

3. Lee DW, Santomaso BD, Locke FL, et al. ASTCT consensus grading for cytokine release syndrome and neurologic toxicity associated with immune effector cells. *Biol Blood Marrow Transplant.* 2019;25:625-638.



Completed Research: PRACTICE MANAGEMENT RESEARCH\

Evaluation of a Decentralized Infusion Pharmacist in an Oncology Infusion Clinic of an Academic Medical Center

Presenting Author: Sarah Castelo, PharmD, BCOP, Vanderbilt University Medical Center, Nashville, TN

Co-Authors Nick Hopkins, PharmD, BCSCP, Vanderbilt University Medical Center, Nashville, TN

BACKGROUND: Pharmacists trained in oncology oversee and manage the use of oncology medications from a core pharmacy within the oncology infusion center. There are limited data on the value of a decentralized infusion pharmacist in an oncology infusion center.

OBJECTIVES: To improve efficiency of infusion clinic workflow by integrating an oncology-trained infusion pharmacist with the nursing staff to assist with troubleshooting, problem solving, and communication with the core pharmacy and the clinical pharmacy specialists. Also, to improve patient care by providing more direct access to a pharmacist.

METHODS: This quality improvement study was conducted at an oncology infusion center at Vanderbilt University Medical Center, Nashville, TN, an academic medical center . A pharmacist worked in the decentralized infusion pharmacist role during peak clinic hours when the operations of the pharmacy could be safely and efficiently managed without the pharmacist physically located inside the core pharmacy. The decentralized infusion pharmacist worked in the oncology infusion clinic, integrated with the nursing staff. Data collected included the number of bedside double-checks for chemotherapy conducted, the number of questions fielded from nursing and the core pharmacy, communication with clinical pharmacy specialists, and patient education. The authors conducted an online satisfaction questionnaire after the pilot program was complete.

RESULTS: A total of 1796 patient visits were completed on 20 separate days from January 2024 to August 2024, when decentralized infusion pharmacist coverage was provided. The most common tasks performed by the decentralized infusion pharmacist included fielding questions

from nursing (n=209), bedside double-checks for chemotherapy (n=156), and fielding questions from the core pharmacy (n=86). Of the staff members who completed the satisfaction questionnaire, 60% of nurses reported feeling less busy with the integration of a decentralized infusion pharmacist; whereas 54% of pharmacists in the core pharmacy reported no significant difference in their workload.

DISCUSSION: Decentralization of pharmacy services includes moving the decision-making closer to users. By adapting departmental workflow, pharmacy services are better able to support nurses. There was continuous effort to refine processes to address nursing and pharmacy concerns.

CONCLUSIONS: The integration of a decentralized infusion pharmacist in the oncology infusion clinic improved accessibility to a pharmacist, improved clinic workflow for the nursing staff, and increased the ability to troubleshoot issues. Further studies are needed to fully evaluate the effects the decentralized infusion pharmacist role would have on patient-related outcomes.

1. Baroody C, Sandler M, Hong C, et al. Evaluation of a decentralized investigational drug service pharmacist in a cancer clinical trial infusion unit. *J Oncol Pharm Pract.* 2024;30:1200-1206.
2. Chacko R, Worsham T, Gass J. Assessment of nursing satisfaction with pharmacy services. Poster presented at: ASHP 2019; December 8-12, 2019.
3. Mashni OK, Nazer LH, Khalil, HZ, et al. Impact of clinical pharmacy services on patient management in the chemotherapy infusion clinics: a 5-year study at a comprehensive cancer center. *J Pharm Pract.* 2022;35:686-690.

Completed Research: PRACTICE MANAGEMENT RESEARCH

Evaluation of Reduced Observation Times for Subcutaneous Administration of Daratumumab and Hyaluronidase in Patients With Multiple Myeloma

Presenting Authors: Arsemen Brown, PharmD, Alisha Desai, PharmD, Elizabeth Graver, PharmD, Amy Zheng, PharmD, Sandra Horowitz, PharmD, and Brian Primeaux, PharmD, BCOP, MD Anderson Cancer Center, Houston, Texas

BACKGROUND: Daratumumab is an anti-CD38 monoclonal antibody that can be given subcutaneously(SC) when formulated with hyaluronidase for previously untreated and relapsed/refractory multiple myeloma. Infusion-related reactions are often observed with daratumumab, and premedication is required before administration. However, the manufacturer provides little guidance on observation times and monitoring after daratumumab/hyaluronidase SC administration. Our institution originally adopted observation times similar to research protocols conducted within the institution of 6 hours for the first dose, 2 hours for the second dose, and 15 minutes for subsequent doses of daratumumab/hyaluronidase SC. Since then, publications have demonstrated the appropriateness of shorter monitoring periods of varying degrees. Due to these data, observation times after daratumumab/hyaluronidase SC administration were reduced to 2 hours for the first dose, 30 minutes for the second dose, and 15 minutes for subsequent doses.

OBJECTIVES: The primary objective was to determine if reduced observation times for daratumumab/hyaluronidase SC affected the rates of infusion reactions. The secondary objective was calculation of chair time saved.

METHODS: All daratumumab/hyaluronidase SC administrations during a 2-year timeframe (February 4, 2021, to May 1, 2023) were collected via Pharmacy Analytics. Administrations were separated based on preintervention and postintervention. Preintervention was defined as SC daratumumab administrations with an observation time of 6 hours after the first dose and 2 hours after the second dose. Postintervention was defined as SC daratumumab administrations with an observation time of 2 hours after the first dose and 30 minutes after the second dose.

RESULTS: A total of 203 patients were included, with 87 in the preintervention group and 116 in the postintervention group. No clinically significant differences were found in rates of reaction with reduced observation time with daratumumab/hyaluronidase SC. Reaction were similar, with 6% for dose 1 and 1% for dose 2 for preintervention group, and 4% for dose 1 and 3% for dose 2 in the postintervention group. The most common reactions were hypertension followed by shortness of breath. For the 116 patients in the postintervention group, reduced observation time saved nearly 27 days of chair time.

CONCLUSION: The results of this study show there was no clinically significant difference in infusion related reactions when comparing longer observation times to shortened observation times. Shortened observation times resulted in a significant amount of chair time saved. Based on the results of this study, it is recommended to continue the shortened observation time after administering daratumumab/hyaluronidase SC.

1. Darzalex Faspro (daratumumab and hyaluronidase-fihj) injection, for subcutaneous use [prescribing information]. Janssen Biotech, Inc.; July 2024. *[Accessed 5 Sept 2024]* www.accessdata.fda.gov/drugsatfda_docs/label/2024/761145s0251bl.pdf
2. Mateos MV, Nahi H, Legiec W, et al. Subcutaneous versus intravenous daratumumab in patients with relapsed or refractory multiple myeloma (COLUMBA): a multicentre, open-label, non-inferiority, randomised, phase 3 trial. *Lancet Haematol.* 2020;7:e370-e380. Erratum: *Lancet Haematol.* 2020;7:e710.
3. Tam AH, Jung Y, Young R, et al. Evaluation of subcutaneous daratumumab injections in the ambulatory care setting. *J Oncol Pharm Pract.* 2022;28:1819-1825.
4. Maples KT, Hall KH, Joseph NS, et al. Eliminating the monitoring period with subcutaneous daratumumab: a single-center experience. *Blood Cancer J.* 2023;13:29.
5. Hughes DM, Henshaw L, Blevins F, et al. Standard 30-minute monitoring time and less intensive pre-medications is safe in patients treated with subcutaneous daratumumab for multiple myeloma and light chain amyloidosis. *Clin Lymphoma Myeloma Leuk.* 2022;22:566-568.

6. Davis JA, Youngberg H, Gaffney K, et al. 'Fast but not so furious': short observation time after subcutaneous daratumumab administration is both a safe and cost-effective strategy. *Clin Lymphoma Myeloma Leuk.* 2022;22:e680-e684.



Completed Research: PRACTICE MANAGEMENT RESEARCH

Evaluation of Services Provided by Pharmacists in Addition to Accredited Patient Management Program Responsibilities in a Health-System Specialty Pharmacy

Presenting Authors: Maria Pinto, PharmD, University of Rochester Community Pharmacy, Rochester, NY; Elizabeth Rightmier, PharmD, BCPS, University of Rochester Specialty Pharmacy, Rochester, NY; Jeremiah Moore, PharmD, BCOP, James P. Wilmot Cancer Institute, University of Rochester Specialty Pharmacy

BACKGROUND: Health-system specialty pharmacies struggle to maintain adequate staffing with continuous growth. New positions are typically justified by increased prescription volume; however, pharmacists are frequently tasked with providing services for patients not directly falling under responsibilities of their accredited patient management program. Guidance on services and time spent may be a valuable tool for health-system specialty pharmacies to consider in requesting additional support.

OBJECTIVE: To approximate the quantity and type of pharmacist actions related to patients fulfilling their specialty prescriptions at external pharmacies or nonspecialty medications.

METHODS: Single-center, prospective quality improvement study to evaluate pharmacy services requested for patients either not actively enrolled in a University of Rochester Specialty Pharmacy (URSP) patient management program or related to medications fulfilled externally. URSP is accredited by the Utilization Review Accreditation Commission and the Accreditation Commission for Healthcare. External was defined as any pharmacy other than URSP, which included ambulatory, infusion, or inpatient pharmacy sites within the health-system. Standardized actions performed by clinical pharmacists employed by the specialty pharmacy and with integrated workflows in the nononcology (cardiology, dermatology, gastroenterology, hepatology, infectious diseases, neurology, primary care, rheumatology) and oncology clinics were documented in the electronic medical record between September 1, 2023 and November 30, 2023. Primary outcome was quantity of pharmacist actions requested to intervene with medication therapy problems or to answer drug information questions. Secondary outcomes included source of the request for pharmacy services, drug category (specialty, nonspecialty,

infusion), status of enrollment in the patient management program, urgency of request, and approximate time spent. All analysis was descriptive.

RESULTS: Pharmacists performed 926 actions, documenting 775 encounters involving 630 unique patients. Majority of pharmacy service requests were from the oncology clinic (76.3%), followed by rheumatology (11.2%) and neurology (5%). Among oncology clinic, providers most requested pharmacy services (57.5%), specifically advanced practice providers (39.6%) and physicians (17.9%). Most actions were non-urgent, representing 68.9% for oncology and 84.8% for non-oncology clinics. Actions were most frequently categorized as drug information (68.9%) followed by medication therapy problems (16.5%). The most common pharmacy service requests from oncology clinics involved nonspecialty medications (47.8%) and infusions (30.7%); whereas nononcology clinics primarily received requests related to specialty medications fulfilled externally (61.6%). Pharmacists spent <15 minutes (68.8%) and 15 to 30 minutes (24.1%) per patient on most encounters.

CONCLUSION: This study highlights the significant contributions that URSP provides to patients in addition to their responsibilities within the scope of their specialty pharmacy–accredited patient management programs and supports the request for additional pharmacist support.

1. Tichy EM, Hoffman JM, Tadrous M, et al. National trends in prescription drug expenditures and projections for 2023. *Am J Health Syst Pharm.* 2023;80:899-913.
2. West S, Filstein D. Outlook for health system-based specialty pharmacy in 2022 and beyond. *Pharm Times.* January 20, 2022. www.pharmacytimes.com/view/outlook-for-health-system-based-specialty-pharmacy-in-2022-and-beyond
3. Zuckerman AD, Mourani J, Smith A, et al. 2022 ASHP survey of health-system specialty pharmacy practice: clinical services. *Am J Health Syst Pharm.* 2023;80:827-841.
4. Cobaugh DJ. Health-system specialty pharmacy services: setting the standard. *Am J Health Syst Pharm.* 2023;80:787-788.

5. Proceedings of the ASHP Specialty Pharmacy State of Practice in Hospitals and Health Systems-Future Directions Summit. *Am J Health Syst Pharm*. 2021;78:1800-1823.
6. James D. Specialty lite model goes beyond improving adherence. Pharmacy Times. April 30, 2019. www.pharmacytimes.com/view/specialty-lite-model-goes-beyond-improving-adherence
7. Armistead LT, Cruz A, Levitt JM, et al. Medication therapy problem documentation and reporting: a review of the literature and recommendations. *J Am Coll Clin Pharm*. 2021;4:1337-1347.
8. Haas CE, Dick TB. Productivity, workload, and clinical pharmacists: definitions matter. *Am J Health Syst Pharm*. 2022;79:728-9.
9. Rough SS, McDaniel M, Rinehart JR. Effective use of workload and productivity monitoring tools in health-system pharmacy, part 1. *Am J Health Syst Pharm*. 2010;67:300-311.
10. Rough SS, McDaniel M, Rinehart JR. Effective use of workload and productivity monitoring tools in health-system pharmacy, part 2. *Am J Health Syst Pharm*. 2010;67:380-388.

HOPAS

Together we can reach new heights

Completed Research: PRACTICE MANAGEMENT RESEARCH

Evaluating the Impact of Interactive Ambulatory Care Discussions at an Oncology Center

Presenting Authors: Abigail Shockley, PharmD, BCOP, JP Sanchez, PharmD, BCOP, and Todd W. Canada, PharmD, BCNSP, BCCCP, FASHP, FTSH, FASPEN, Division of Pharmacy, University of Texas MD Anderson Cancer Center, Houston, TX

BACKGROUND: An interactive Ambulatory Shared Pharmacy Student Instruction (ASPSI) program was developed to supplement ambulatory care clinic rotations at a large oncology academic medical center. However, traditional ambulatory care disease management is mostly absent due to the specialized nature of these oncology clinics. Pharmacy students completing advanced pharmacy practice experiences (APPE) primarily engage in specialized oncology-related patient care during their rotations, which limits their ability to fully apply didactic knowledge and gain practical application of common ambulatory care diseases such as diabetes mellitus or hypertension during their 6-week rotation.

OBJECTIVE: To evaluate the perceptions of 4th year APPE students on their ambulatory care disease expertise after participation in ASPSI.

METHODS: ASPSI is a series of case-based interactive topic discussions to enhance learning on designated topics for APPE students during their ambulatory care rotation(s) at MD Anderson Cancer Center. Four discussions are held during each 6-week rotation. The discussions are diabetes mellitus, hypertension, constipation/diarrhea, and anticoagulation. Discussions are intended to be interactive learning experiences while led and facilitated by ambulatory care oncology clinical pharmacy specialists. Over the past academic year, each student involved in these topic discussions was asked to voluntarily complete an anonymous survey at the end of the 6 weeks. Students were asked to rate their confidence level on a 5-point Likert-type scale for the ability to care for a patient on a scale of 1 to 5, with 1 being very low confidence and 5 being very high confidence. Survey results were compiled and described.

RESULTS: All students (N=11) completed the survey, with 100% reporting these discussions improved their confidence in managing patients with diabetes mellitus, hypertension,

constipation/diarrhea, and anticoagulation. The mean confidence score for diabetes management increased from 2.36 to 3.55, for hypertension from 2.82 to 4.18, for anticoagulation from 2.82 to 4.09, and for constipation/diarrhea from 2.45 to 4.18. Median confidence scores across all conditions also showed a shift from 2 to 3 before the discussions to 4 after the discussions. These results indicate a marked increase in student confidence in managing these conditions after participating in the interactive discussions.

CONCLUSIONS: Student evaluations of the interactive ASPSI topic discussions were overwhelmingly positive in a nontraditional ambulatory care APPE setting. The results support the continued use of this instructional approach to reinforce core ambulatory care topics during specialized oncology APPE ambulatory care rotations. Ongoing use of student feedback through surveys will allow preceptors to improve the program and potentially expand the range of topics covered in future iterations.

1. Sullivan GM, Artino Jr AR. Analyzing and interpreting data from likert-type scales. *J Grad Med Educ.* 2013;5:541-542.

HOPAC

Together we can reach new heights

Completed Research: PRACTICE MANAGEMENT RESEARCH

Hematologic Oral Oncolytics: Adherence, Healthcare Cost and Utilization

Presenting Authors: Francis C. Staskon, PhD, Khadijah Khan, CPhT, and Laly Havern, PharmD, Walgreen Co., Deerfield, IL; Amy Pfeifer, PharmD, BCPS, CSP, Walgreen Co., Appleton, WI

BACKGROUND: Connected Care Oncology is a patient-centered clinical program that includes a set of 21 unique oral hematologic oncolytic agents for affected oncology patients.


OBJECTIVE: To identify significant associations between a discontinuation adherence metric for the set of Connected Care Oncology hematologic oral oncolytic agents and total medical costs, hospitalizations, and total length of stay (LOS) for inpatients. Research was deemed exempt from HIPAA (Health Insurance Portability and Accountability Act) by a Walgreens Advarra Independent Review Board #39505.

METHODS: A retrospective cohort design of patients was used from the MarketScan Commercial Claims and Encounters for 2022. The sample selection required ≥ 2 fills of targeted medication from the 2022 files with a primary cancer diagnosis code in medical files for 2021 or 2022, and patients must have been continuously enrolled and aged 18 to 64 years. Exclusion criteria were presence of hospice care or organ transplants and those starting medication therapy in the last 45 days of 2022. Discontinuation was indicated by a gap exceeding 1.5 times (previous days' supply) on consecutive fills. General linear models predicted total medical costs, hospitalizations, and LOS (with gamma or logit links). Predictors included discontinuation, and 12 covariates (including COVID-19 indications, surgeries, Charlson Comorbidity Index comorbidities [less cancer], combination therapy, inpatient and outpatient utilization levels, demographics, and insurance type) and their interaction terms. The economic valuation compared the model predictions by discontinuation status.

RESULTS: A total of 4296 patients in 2022 met sample and model criteria, with 72.3% adherent. Predicted total medical costs significantly increased for nonadherent patients compared with adherent patients (\$73,663 per member per year [PMPY]; $P < .0001$). When considering the additional costs associated with pharmacy adherence, the predicted total medical and pharmacy

costs remained increased for nonadherent patients by (\$19,193 PMPY; $P<.46$). Compared with adherent patients, odds of hospitalization were significantly increased ($OR=2.22$; $P<.0001$) and LOS was significantly longer for nonadherent patients (4 days; $P<.0001$). When considering the difference in hospital admissions rate and LOS when admitted, the predicted inpatient medical spend was higher for the nonadherent patients (\$22,691 PMPY). Details on all significant model covariates and outcomes as well as economic valuation comparisons are presented in poster.

CONCLUSION: Being adherent to oral hematologic oncolytics can lead to lower medical costs, odds of hospitalization, and LOS after controlling for many other influences on these outcomes.



HOPAC

Together we can reach new heights

Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

High Antimicrobial Resistance in a Pediatric Oncology Unit in Malawi Underscores Relevance of Stronger Antimicrobial Stewardship in Sub-Saharan Africa

Presenting Authors: Agness Chisomo Chitedze,MSc,BPharm Steven Kafora Mlenga,BSc and Rizine Mzikamanda,MBBS Baylor College of Medicine Children's Foundation Malawi; Tarsizio Maida,MSc Ministry of Health, Kamuzu Central Hospital, Lilongwe, Malawi; Casey McAtee,MD,MPH and Nmazuo Ozuah,MD Texas Children's Global HOPE Program, Houston Texas, and Department of Pediatrics, Section of Hematology-Oncology, Baylor College of Medicine, Houston Texas

BACKGROUND: Antimicrobial resistance (AMR) is a global threat contributing to significant deaths worldwide. Pediatric patients with cancer are prone to serious infections requiring antimicrobial treatment. There is paucity of data on AMR in pediatric oncology units in sub-Saharan Africa, where routine blood cultures are often unavailable.

OBJECTIVE: To describe the prevalence of AMR in a treatment center in Malawi to inform the need for stronger advocacy for antimicrobial stewardship in the region.

METHODS: This was a retrospective cross-sectional study of blood cultures obtained between January 2019 and August 2023 from pediatric oncology patients at Kamuzu Central Hospital in Lilongwe, Malawi. Multidrug resistance was defined as lack of susceptibility to ≥ 1 antimicrobial drug in ≥ 3 antimicrobial categories. Association between categorical variables was measured with Pearson's chi-square test, and multivariate logistic regression was used to test for association between multiple variables.

RESULTS: We identified 447 blood culture results from 262 patients. Most were obtained from patients with leukemia (37.5%), lymphoma (28.6%), or solid tumors (21.4%). The blood culture positivity was 37.9% (169/447). Of these, 32.5% (55/169) were identified as contaminants. Of the remaining 114 true isolates, 70.2% were gram-negative (50% Enterobacteriaceae, with *Klebsiella* as the most common). The prevalence of multidrug resistance was 54.4% (95% confidence interval, 44.8%-63.7%), with rates of 68% versus 41% for gram-negative and -positive organisms, respectively ($P=.04$). Of the gram-negative isolates with sensitivity data,

resistance was as follows: gentamicin (68%), ceftriaxone (67%), ciprofloxacin (59%), ceftazidime (52%), imipenem (47%), and piperacillin-tazobactam (38%). No significant associations were found between multidrug resistance rate and cancer diagnosis.

CONCLUSION: The prevalence of AMR in our population is higher than described in developed countries, emphasizing critical need for routine blood cultures in pediatric oncology units as a strategy for surveillance and to aid antimicrobial stewardship. High contamination rates denote the need for implementing proper blood culture collection techniques.

1. Antimicrobial stewardship programmes in health-care facilities in low- and middle-income countries: a WHO practical toolkit. World Health Organization. October 22, 2019. Accessed Date May 2024 www.who.int/publications/i/item/9789241515481
2. Muratore E, Baccelli F, Leardini D, et al. Antimicrobial stewardship interventions in pediatric oncology: a systematic review. *J Clin Med*. 2022;11:4545.
3. Papan C, Reifenrath K, Last K, et al. Antimicrobial use in pediatric oncology and hematology in Germany and Austria, 2020/2021: a cross-sectional, multi-center point-prevalence study with a multi-step qualitative adjudication process. *Lancet Reg Health Eur*. 2023;28:100599.

Together we can reach new heights

Completed Research: PRACTICE MANAGEMENT RESEARCH

Identifying Barriers to Expanded Access Program Patient Enrollment in the Hematology/Oncology Community Surveying Pharmacists

Presenting Authors: Jahmal Williams, PharmD, RPh, Claire Groce, PharmD, Madison Henry, PharmD, RPh, Kyle Taylor, PharmD, RPh, Nik Mohan, PharmD, Joseph Barone, PharmD, FCCP, Michael Toscani, PharmD, Ernest Mario School of Pharmacy, Piscataway, NJ; Paul Weber, MD, RPh, MBA, Ernest Mario School of Pharmacy and Rutgers Robert Wood Johnson Medical School, Piscataway, NJ

BACKGROUND: Expanded access programs (EAPs) provide a pathway for patients with serious or life-threatening conditions to obtain investigational treatments outside of clinical trials, when no satisfactory or comparable treatment options exist, or they do not qualify for a clinical trial. Use is limited by a number of challenges, including poor access to information, document burden, and lack of response from drug sponsors regarding ability to supply drug.¹ This study aims to gauge the baseline knowledge gaps and barriers that may deter oncology/hematology pharmacists from enrolling patients into an EAP.

OBJECTIVES: To assess the knowledge of US academic and community oncology/hematology pharmacists regarding EAPs and identify barriers to patient access through these programs

METHODS: Dissemination of a Rutgers electronic Institutional Review approved, protocol #Pro2024000713 voluntary Qualtrics survey through the Hematology/Oncology Pharmacy Association's email list. The research survey was sent only to US members who practice clinical oncology/hematology pharmacy in academic and/or community settings. Students, residents, fellows, technicians, and pharmaceutical industry members were excluded from the study. The survey was disseminated in June 2024 and was open until the end of July 2024.

RESULTS: After reviewing 38 survey responses, our findings suggest pharmacists working in an academic setting were more aware of EAPs than those working in a community setting. A total of 69% of participants requested use of a drug through an EAP. However, most (63%) participants were slightly knowledgeable or not knowledgeable at all when it comes to finding

access to investigational drugs for patients via EAPs. The information sources varied among pharmacists, with the most common being the pharmaceutical manufacturer (59%). The top barriers included time to enroll patients (66%), documentation burden (59%), and lack of familiarity with EAP processes (56%).

CONCLUSIONS: Pharmacists, as part of the interprofessional care team, play a critical role in decision-making during a patient's treatment journey. EAPs are an option for patients who have limited or no treatment options remaining. This study identifies key knowledge gaps and barriers to treatment access via EAPs. These results provide an opportunity to close knowledge gaps and barriers related to EAPs by providing continuing education credits to pharmacists in the academic and community settings. Informing oncology/hematology pharmacists in academic and community centers regarding EAPs allows the opportunity to educate and offer alternative treatment options to patients, providing hope to patients and families. Further research will be needed to assess the unmet educational needs and the potential benefits of increased education on patient care and outcomes.

1. Fu S, Muquith M, Williams EF, et al. Barriers to implementing expanded access programs in oncology: a multidisciplinary perspective. *J Clin Oncol*. 2023;41(16 suppl):e18632.

Together we can reach new heights

Completed Research: PRACTICE MANAGEMENT RESEARCH

Impact of Clinically Embedded Oncology Pharmacists on Immune Checkpoint Inhibitor Outcomes

Presenting Authors: Colton Zwart, PharmD, BCOP, and Mark Wagner, PharmD, BCOP, Cowell Family Cancer Center, Munson Healthcare, Traverse City, MI; Karen Farris, PhD, University of Michigan College of Pharmacy, Ann Arbor, MI; David Reyes-Gastelum, and Katie Young, Michigan Institute for Care Management and Transformation, Ann Arbor, MI; Emily Mackler, PharmD, BCOP, Michigan Oncology Quality Consortium, Michigan Institute for Care Management and Transformation, YesRx, and University of Michigan, Ann Arbor, MI

BACKGROUND: Multiple publications have highlighted the pharmacist's role in managing immune-related adverse events (AEs) from immune checkpoint inhibitor (ICI) therapy for patients with cancer.

OBJECTIVE: The purpose of this study was to compare outcomes in patients initiating ICI therapy for cancer before and after the integration of 2 clinically embedded clinical pharmacists in a large community oncology practice via the Pharmacists Optimizing Oncology Care Excellence in Michigan program.

METHODS: This single-center, retrospective analysis was conducted with data from March 2018 to March 2024. The intervention was clinically embedded pharmacist care, which was identified by a pharmacist education visit. Up to 134 patients in each of the pre- and postintervention groups were identified. Charts were eligible for abstraction if the patients were newly prescribed an ICI for cancer. The following data were abstracted: patient demographics, ICI medication, therapy intent (curative vs palliative), cancer diagnosis, start date, education date, corticosteroid use, follow-up visits, emergency department and hospitalizations, reason for emergency department/hospitalization, and reasons for ICI discontinuation. Chi-square analysis compared the pre- and postintervention time periods.

RESULTS: Abstraction was completed for 133 patients in the preintervention group and 134 patients in the postintervention group. The demographics in the preintervention group versus the

postintervention group included a mean age of 67 years (standard deviation [SD], 12.3) versus 69 years (SD, 11.5), respectively; 64% female versus 70% female, respectively; 96% white versus 97% white, respectively; and 17% curative versus 34% curative, respectively. The most common cancer diagnoses were lung, endometrial, melanoma, kidney, breast, and bladder. Pembrolizumab was utilized most frequently in 43% of the preintervention patients and 52% of the postintervention patients. More patients in the postintervention group received education before starting ICI therapy (99% vs 86%; $P=.002$). The postintervention group had a 41% discontinuation rate due to AEs, progression, or hospice/death compared with 51% in the preintervention group ($P=.098$). Steroid use was similar between groups, with 27% of patients in the preintervention group requiring steroids and 25% in the postintervention group. There were no differences between the groups in the time to steroid initiation or the time receiving a steroid. The postintervention group had less hospitalizations (24% vs 41%; $P=.009$) and less emergency department visits (28% vs 46%; $P=.005$) within the first 6 months of therapy for cancer or treatment related AEs compared with the preintervention group.

CONCLUSION: Embedded clinical pharmacist care improved the outcomes of patients receiving ICIs for the treatment of cancer, including improved rates of education and decreased incidence of unplanned healthcare utilization.

1. Le S, Chang B, Pharm A, Chan A. Impact of pharmacist-managed immune checkpoint inhibitor toxicities. *J Oncol Pharm Pract.* 2021;27:596-600.
2. Renna CE, Dow EN, Bergsbaken JJ, Leal TA. Expansion of pharmacist clinical services to optimize the management of immune checkpoint inhibitor toxicities. *J Oncol Pharm Pract.* 2019;25:954-960.
3. Myers G, Stevens J, Flewelling A, et al. Evaluation and clinical impact of a pharmacist-led, interdisciplinary service focusing on education, monitoring and toxicity management of immune checkpoint inhibitors. *J Oncol Pharm Pract.* 2023;29:145-154.
4. POEM program. Accessed October 4, 2024. www.moqc.org/poem

Completed Research: PRACTICE MANAGEMENT RESEARCH

Impact of Clinically Embedded Pharmacists on the Time of Other Oncology Clinicians

Presenting Authors: Emily Mackler, PharmD, BCOP, Michigan Oncology Quality Consortium, Michigan Institute for Care Management and Transformation, YesRx, and University of Michigan College of Pharmacy, Ann Arbor, MI; Maisa Nazzal, PharmD, MS, and Karen Farris, PhD, University of Michigan College of Pharmacy, Ann Arbor, MI; Maja Hill, PharmD, Lemon-Holton Cancer Pavilion, Corewell Health, Grand Rapids, MI; Emily Johengen, PharmD, BCACP, Trinity Health IHA Hematology Oncology, Ypsilanti, MI; Katie Sias, PharmD, BCOP, Pardee Cancer Wellness Center, MyMichigan Health, Midland, MI; Samuel Snowaert, PharmD, BCOP, MBA, Covenant Cancer Care Center, Saginaw, MI; Jennifer VanSickler, PharmD, Herbert-Herman Cancer Center, University of Michigan Health - Sparrow, Lansing, MI; Mark Wagner, PharmD, BCOP, and Colton Zwart, PharmD, BCOP, Cowell Family Cancer Center, Munson Healthcare, Traverse City, MI; Katie Young and David Reyes-Gastelum, Michigan Institute for Care Management and Transformation, Ann Arbor, MI

BACKGROUND: Clinical pharmacists save physicians' time through interventions and clinical services offered in the emergency department, but studies evaluating time savings in oncology practice have not been done.¹ Clinical pharmacists embedded in ambulatory care practices provide a variety of services, such as selecting appropriate chemotherapy regimens and supportive care, reviewing medication lists and laboratory tests, managing adverse events, educating patients, and answering drug information questions. Administratively, they may also build order sets, provide staff education, and release orders. The purpose of this study was to describe the time spent by embedded clinical oncology pharmacists and who that effort would be attributed to if they were not in the clinic.

METHODS: On 2 different weeks, 7 pharmacists from Pharmacists Optimizing Oncology Excellence in Michigan practices recorded their activities using a data collection form that included patient and nonpatient focused activities.² The second data collection is the basis for this analysis, and data were collected in July 2024. The 7 pharmacists provided 922 entries grouped into administration, chart review/release orders, care coordination, clinical question,

documentation, follow-up, order management, teach, staff Education, and other. The number of minutes attributed to the activity and the professional who would do the activity if the clinical pharmacist was not present were recorded. Hours per activity and the substituted professional were determined.

RESULTS: Pharmacists recorded activities for an average of 42 hours per week, and 70% of activities were patient focused. Among patient focused activities, pharmacists' distribution of time was primarily focused on chart reviewing and releasing orders, teaching, and documentation. For patient-focused activities, pharmacists substituted a mean of 5 hours for advance practice practitioners, 3.3 hours for physicians, 1 hour for medical assistants, 11.8 hours for nurses, 4.1 hours for another pharmacist, and 3.7 hours for no one in the clinic on a per pharmacist per week basis. For nonpatient focused activities, the clinical pharmacists substituted 7.3 hours for no one in the clinic and 4.1 hours for another pharmacist, per pharmacist per week.

CONCLUSION: Oncology pharmacists' time was 70% patient focused, and they saved on average 20 hours per week per pharmacist for nurses, advance practice practitioners, and physicians.

1. Grill J, Bryant C, Markel K, Wisniewski SJ. A study of time saved by emergency medicine physicians through working with clinical pharmacists in the emergency department. *Am J Emerg Med.* 2019;37:1720-1722.
2. POEM program. Accessed October 4, 2024. www.moqc.org/poem

Together we can reach new heights

Completed Research: PRACTICE MANAGEMENT RESEARCH

Impact of Embedded Clinical Pharmacists on Health Utilization Outcomes in a Large Community Oncology Practice

Presenting Authors: Mark Wagner, PharmD, BCOP, and Colton Zwart, PharmD, BCOP, Cowell Family Cancer Center, Munson Healthcare, Traverse City, MI; Karen Farris, PhD, University of Michigan College of Pharmacy, Ann Arbor, MI; David Reyes-Gastelum and Katie Young, Michigan Institute for Care Management and Transformation, Ann Arbor, MI; Emily Mackler, PharmD, BCOP, Michigan Oncology Quality Consortium, Michigan Institute for Care Management and Transformation, YesRx, and University of Michigan College of Pharmacy, Ann Arbor, MI

BACKGROUND: Pharmacists play a critical role in team-based care, and this is well established in the primary care setting.

OBJECTIVE: To compare healthcare utilization outcomes before and after the integration of 2 embedded clinical pharmacists in a large community oncology practice via the program, Pharmacists Optimizing Oncology Excellence in Michigan.

METHODS: This single-center, retrospective analysis was conducted with data from January 2018 to March 2024. The intervention was clinically embedded pharmacist care, identified by a pharmacist education visit. Up to 400 patients in each of the pre- and postintervention groups were identified. Charts were eligible for abstraction if patients were newly prescribed anticancer therapy for any of the 3 following areas: gynecologic oncology diagnosis, the use of immune checkpoint inhibitor(s), or the use of an oral anticancer agent(s). The following data were abstracted: patient demographics, medication, cancer diagnosis, emergency department visits and hospitalizations, and reason for emergency department/hospitalization. A chi-square analysis compared pre- and post-intervention time periods.

RESULTS: Abstraction was completed for 398 patients in the preintervention group and 350 patients in the post-intervention group. No demographic differences were detected between groups. The pre-intervention group had a mean age of 67 years (standard deviation (SD), 11.5),

69% were female, and 96% were White. The postintervention group had a mean age of 68 years (SD, 12.5), 57% were female, and 97% were White. The majority of patients in both groups had either ovarian, endometrial, non–small cell lung, prostate, or breast cancer diagnosis. At 6 months following treatment initiation, the pre-intervention group had a 24% incidence of hospitalization due to cancer symptoms, treatment adverse events, or unknown reason versus 19% in the post-intervention group ($P=.065$). Emergency department visits at 6 months for the aforementioned reasons were 32% in the pre-intervention group versus 25% in the post-intervention group ($P=.035$).

CONCLUSION: Embedded clinical pharmacist care reduced the incidence of emergency department visits in the first 6 months of anticancer treatment initiation.

1. Coe AB, Choe HM, Diez HL, et al. Pharmacists providing care in statewide physician organizations: findings from the Michigan Pharmacists Transforming Care and Quality Collaborative. *J Manag Care Spec Pharm*. 2020;26:1558-1566.
2. POEM program. Accessed October 4, 2024. www.moqc.org/poem

HOPANA

Together we can reach new heights

Completed Research: PRACTICE MANAGEMENT RESEARCH

Improving Oral Anticancer Medication Adherence/Adverse Event Assessment Documentation

Presenting Authors: Grace Morrison, BS, and Lisa Holle, PharmD, BCOP, FHOPA, FISOPP, University of Connecticut, Storrs, CT; Andrea Moran, APRN-BC, DNP, and Margaret Savage, BSN, RN, UConn Health Carole & Ray Neag Comprehensive Cancer Center; Michael K. Keng, MD, University of Virginia

BACKGROUND: To improve patient safety, most oral anticancer medication (OAM) programs monitor adverse events (AEs) and adherence within 7 to 14 days following OAM initiation. At our institution, we identified key tasks associated with documenting OAM adherence and AEs during cycle 1. However, only 50% of these tasks were appropriately documented.

OBJECTIVE: As part of the 2022-2023 Hematology/Oncology Pharmacy Association/American Society of Clinical Oncology quality training program, we conducted a quality improvement project aimed at improving our task documentation rate to >75%.

METHODS: We engaged a multidisciplinary team to assist in gathering and interpreting data. A process map was created, and barriers were identified. A cause-and-effect diagram, Pareto chart, and action prioritization matrix were used to narrow the focus of improvement changes initially and after each cycle. Three plan-do-study-act (PDSA) cycles were conducted (January 2023 to January 2024). Outcome measure was documentation of AEs/adherence by day 10 of cycle. Our study included all patients with cancer initiating OAM and excluded those on intravenous/oral regimens and gynecology/oncology patients. Process measure was percentage of 9 crucial tasks identified as necessary to appropriately document AEs/adherence. Balance measures included provider visits, emergency department visits, and telephone/electronic medical record OAM-related messages.

RESULTS: PDSA cycle 1 (education on documenting patient education) was implemented in January 2023. PDSA cycle 2 (retraining nurses on OAM AEs/adherence documentation process) was implemented in March 2023. PDSA cycle 3 (standardized cycle 1 OAM process and created

a tool to facilitate communication of OAM start date) was implemented in September 2023. Improvement in our outcome measure (percentage of 9 essential tasks for documenting OAM adherence/AEs in cycle 1) was improved following PDSA cycles 1, 2, and 3 (50%-67%; $P<.001$). This rate was sustainable in the 2 months following project completion. OAM-related patient provider visits, emergency department visits, and OAM-related telephone messages (ie, balance measures) were decreased ($P<.001$). An improvement in documentation of OAM treatment start date was observed with each additional cycle (10%, 23%, and 83%, respectively) and improvement in AE documentation was similarly observed (10%, 12%, and 83%, respectively).

CONCLUSION: Our results were likely affected by significant nursing turnover, lack of formal OAM process training, and institutional buy-in for OAM program/training. Our standardizing process initiative improved the percentage of AEs/adherence documentation tasks necessary to ensure completion of appropriate OAM monitoring by day 10 of cycle 1 following prescription. This also resulted in fewer OAM-related telephone messages, which not only decreased provider time but signified that fewer potential problems occurred with appropriate AEs/adherence checks.


1. Muluneh B, Schneider M, Faso A, et al. Improved adherence rates and clinical outcomes of an integrated, closed-loop, pharmacist-led oral chemotherapy management program. *J Oncol Pract.* 2018;14:e324-e334.
2. Kav S, Schulmeister L, Nirenberg A, et al. Development of the MASCC Teaching Tool for patients receiving oral agents for cancer. *Supp Care Cancer.* 2010;18:583-590.
3. Betcher J, Dow E, Khera N. Oral chemotherapy in patients with hematological malignancies-care process, pharmacoeconomics and policy implications. *Curr Hematol Malig Rep.* 2016;11:288-294.
4. Enright K, MacMillan M, Lymburner P, et al. Improving documentation of oral chemotherapy at a community cancer center. *J Oncol Pract.* 2015;11:213-215.
5. Neuss MN, Polovich M, McNiff K, et al. 2013 updated American Society of Clinical Oncology/Oncology Nursing Society chemotherapy administration safety standards, including

standards for the safe administration and management of oral chemotherapy. *J Oncol Pract.* 2013;9:5s-13s. Erratum in: *J Oncol Pract.* 2013;9:265.

6. Weingart SN, Brown E, Bach PB, et al. NCCN task force report: oral chemotherapy. *J Natl Comp Cancer Netw.* 2008;6(suppl 3):S1-S14.

7. Mackler E, Segal EM, Muluneh B, et al. 2018 Hematology/Oncology Pharmacist Association best practices for the management of oral oncolytic therapy: pharmacy practice standard. *J Oncol Pract.* 2019;15:e346-e355.

8. Neuss MN, Gilmore TR, Belderson KM, et al. 2016 Updated American Society of Clinical Oncology/Oncology Nursing Society chemotherapy administration safety standards: including standards for pediatric oncology. *Oncol Nurs Forum.* 2017;44:31-43.



Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

Managing Oral Anticancer Therapies in Patients With Early or Advanced HR-Positive HER2-Negative Breast Cancer: A US-based Community Healthcare Provider Survey

Presenting Authors: Jodi L Taraba, PharmD, MS, BCOP, Mayo Clinic, Rochester, MN; Pamela K Ginex, EdD, MPH, RN, OCN, State University of New York at Stony Brook, Stony Brook, NY; Qinli Lily Ma, PhD, Kelli Thoele, PhD, RN, OCN, Alexandra S Vitko, PhD, Astra M Liepa, PharmD, Wambui Gathirua-Mwangi, PhD, Elyse H Panjic, PharmD, and Engels Chou, MS, Eli Lilly and Company, Indianapolis, IN; Hilary Ellis, BA, Charlotte Clewes, MSc, and Joanna de Courcy, BSc, Adelphi Real World, Bollington, UK; Hope S. Rugo, MD, University of California San Francisco Comprehensive Cancer Center, San Francisco, CA

BACKGROUND: In the past decade, the number of options and use of oral anticancer medications (OAMs) has grown among patients with hormone receptor-positive (HR-positive) and HER2-negative breast cancer. Adherence to OAMs is critical to ensure treatment effectiveness and long-term outcomes.¹ However, there is limited evidence on methods used by healthcare providers (HCPs) to support adherence to OAMs.

OBJECTIVE: To describe methods used by HCPs (pharmacists, oncologists, and advanced practice providers [APPs]) at individual and system levels to support adherence to OAMs for HR-positive, HER2-negative breast cancer.

METHODS: Between July 2023 and October 2023, a cross-sectional survey was conducted among HCPs with ≥ 12 months of experience in treating patients with HR-positive, HER2-negative breast cancer in community settings across the United States. A total of 282 HCPs (100 pharmacists, 91 oncologists, and 91 APPs) completed a one-time online questionnaire.

RESULTS: The majority of HCPs (87.0% pharmacists, 95.6% oncologists, and 71.4% APPs) had ≥ 5 years of experience treating patients with breast cancer. Half of HCPs (53.0% of pharmacists, 49.5% of oncologists, and 49.5% of APPs) were extremely or very satisfied with the methods they use in practice to support adherence to OAMs. The 3 most common patient-level methods to support adherence reported by HCPs included dose adjustments (pharmacists: 65.0%, oncologists: 50.5%, and APPs: 79.1%), use of prophylactic medications (pharmacists: 68.0%, oncologists: 39.6%, and APPs: 76.9%), and advising nonpharmacological interventions

(pharmacists: 69.0%, oncologists: 35.2%, and APPs: 60.4%). Retail pharmacists reported higher use of nonpharmacological interventions and lower use of dose adjustments and prophylactic medications than pharmacists working in clinical or specialty settings. Overall, dose adjustment was reported to be most successful across HCPs, with 66.6% of users perceiving this approach as extremely or very successful in supporting adherence; 64.4% and 32.7% reported prophylactic medications and advising nonpharmacological interventions, respectively, to be similarly successful. The most common system-level methods include staff to help navigate insurance approvals (59.9%) and referral to patient assistance programs (59.2%).

CONCLUSION: Dose adjustments and use of prophylactic medications were considered the most successful patient-level methods to support adherence in patients with HR-positive, HER2-negative breast cancer and were reported as the most commonly used by HCPs. Notably, pharmacists and APPs appeared to use all methods more extensively than oncologists, highlighting their vital role in supporting adherence. Broader incorporation of pharmacists into the management of patients receiving OAMs will help enhance the implementation of these methods, with the goal of increasing patient adherence to OAMs.

1. Makubate B, Donnan PT, Dewar JA, et al. Cohort study of adherence to adjuvant endocrine therapy, breast cancer recurrence and mortality. *Br J Cancer*. 2013;108:1515-1524.

Together we can reach new heights

Completed Research: PRACTICE MANAGEMENT RESEARCH

Optimizing Iron Dextran Infusion Protocols to Enhance Efficiency and Patient Throughput in an Outpatient Infusion Center: A Process Improvement Initiative

Presenting Authors: Minhee Kang, PharmD, BCPS, BCOP, and Richard Verstraete, RN, MedStar Georgetown University Hospital, Washington DC

BACKGROUND: Iron dextran 1000 mg is frequently used in our outpatient infusion center to treat iron deficiency anemia,¹ requiring a lengthy 4-hour chair time, including a test dose followed by a 1-hour observation period before administering the full dose.² With increasing patient demand and limited space,³ reducing the total infusion time became critical to enhance efficiency and patient throughput without compromising safety. A formal discussion of the proposal was presented to the infusion pharmacist, the nurse leader of the outpatient infusion center, and the medical director for our anemia management program. All agreed to our proposal.

OBJECTIVES: To reduce the observation period between the iron dextran test dose and full dose from 1 hour to 15 minutes, ultimately reducing overall chair time and improving workflow efficiency.

METHODS: A literature review of iron dextran protocols was conducted, and key stakeholders, including the infusion pharmacist, nurse leader, and medical director for the anemia management program, discussed and approved the proposal. The intervention, which began in May 2024, was implemented in 2 phases:

- Phase 1: 34 patients had a 30-min observation period.
- Phase 2: 50 patients had the observation period reduced to 15 min.

Retrospective data on wait times, chair times, and adverse events were collected through the electronic health record for patients receiving iron dextran from October 2023 to September 2024 to assess pre- and postintervention outcomes.

RESULTS: A total of 119 doses were assessed preintervention, and 84 doses were assessed postintervention. The intervention resulted in a 22.6% reduction in wait time (from 95 to 73.5 min) and a 26.9% reduction in chair time (from 305 to 222.2 min). In all, 4 minor reactions were

reported, with 2 patients discontinuing treatment. All reactions occurred within 10 minutes of the test dose.

The results were not as drastic as expected, primarily due to the need for additional preparation time for the full dose after communication between nurses and pharmacists.

CONCLUSIONS:

Shortening the observation period between iron dextran doses improved infusion center efficiency by reducing chair time and increasing patient throughput, without compromising patient safety. However, further improvement is needed by streamlining the preparation process and enhancing communication between nurses and pharmacists to fully realize the potential of reduced wait times. Future steps include staff education, improving preparation times, and ongoing evaluation of outcomes to ensure sustained and further improvements over the next 6 months.

1. Auerbach M, Wong L, McClintock J, et al. Safety and efficacy of rapid (one hour) single intravenous dose low molecular weight iron dextran for treatment of oral iron intolerant maternal iron deficient anemia. *Blood*. 2015;126:3356.
2. Arastu AH, Elstrott BK, Martens KL, et al. Analysis of adverse events and intravenous iron infusion formulations in adults with and without prior infusion reactions. *JAMA Netw Open*. 2022;5:e224488. Erratum in: *JAMA Netw Open*. 2022;5:e2221567.
3. Wong L, Smith S, Gilstrap M, et al. Safety and efficacy of rapid (1,000 mg in 1 hr) intravenous iron dextran for treatment of maternal iron deficient anemia of pregnancy. *Am J Hematol*. 2016;91:590-593.

Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

Safety and Cost Associated With Fixed-Dose Immunotherapy in Patients With Weight Extremes

Presenting Authors: Cali Lunowa, PharmD, and Christina Davis, PharmD, BCOP, University of Colorado Cancer Center, Aurora, CO; Moriah Cargile, University of Colorado Hospital, Aurora, CO

BACKGROUND: Nivolumab and pembrolizumab were initially approved with weight-based dosing of 3 mg/kg every 2 weeks and 2 mg/kg every 3 weeks, respectively. Later, the FDA approved fixed doses of 240 mg every 2 weeks or 480 mg every 4 weeks for nivolumab and 200 mg every 3 weeks or 400 mg every 6 weeks for pembrolizumab based on pharmacokinetic modeling showing similar areas under the curve compared with weight-based dosages.¹⁻³ However, patients of low and high weights represented a small percentage of the patients included in the simulations,^{1,3} which led to questions regarding safety and cost implications in patients with weight extremes.³

OBJECTIVES: Compare the incidence of adverse events (AEs) in patients weighing ≤ 50 kg or ≥ 110 kg to previously reported literature and determine cost implications of fixed dosing strategies.

METHODS: Retrospective review of adult patients receiving single-agent pembrolizumab or nivolumab for treatment of cutaneous malignancies within the UCHHealth System between January 1, 2016, and May 30, 2024. Patients were included if they received ≥ 1 dose of nivolumab 240 mg or pembrolizumab 200 mg and weighed ≤ 50 kg or > 110 kg. Data collection included baseline demographics, autoimmune disease, line of treatment, earlier treatments, immunotherapy received, dose administered, total number of doses administered per patient, and incidence of AEs. Treatment cost per patient was calculated using standard vial size and present-day vial cost provided from the internal purchasing department. Theoretical cost-savings using weight-based dosing instead of fixed-doses was calculated using the same vial sizes and cost.

RESULTS: A total of 95 patients were assessed, including 45 patients weighing ≤ 50 kg in the low weight group (LWG) and 50 patients weighing ≥ 110 kg in the high-weight group (HWG).

Nivolumab was administered to 67% in the LWG and 56% in the HWG. The mean number of fixed immunotherapy doses per patient was 10 ± 12 for the LWG and 13 ± 14 for the HWG. The observed incidence of AEs with immune checkpoint inhibitors was similar for each group when compared with the literature.³⁻⁵ Fixed doses of nivolumab and pembrolizumab were associated with increased drug cost totaling \$1.8 million when used for patients weighing ≤ 50 kg.

CONCLUSION: Implementation of weight-based nivolumab and pembrolizumab dosing for patients with low weight is a safe and efficient drug-cost saving strategy.

1. Freshwater T, Kondic A, Ahamadi M, et al. Evaluation of dosing strategy for pembrolizumab for oncology indications. *J Immunother Cancer*. 2017;;5:43.
2. Zhao X, Suryawanshi S, Hruska M, et al. Assessment of nivolumab benefit-risk profile of a 240-mg flat dose relative to a 3-mg/kg dosing regimen in patients with advanced tumors. *Ann Oncol*. 2017;28:2002-2008.
3. Long GV, Tykodi SS, Schneider JG, et al. Assessment of nivolumab exposure and clinical safety of 480 mg every 4 weeks flat-dosing schedule in patients with cancer. *Ann Oncol*. 2018;29:2208-2213.
4. Robert C, Long GV, Brady B, et al. Nivolumab in previously untreated melanoma without BRAF mutation. *N Engl J Med*. 2015;372:320-330.
5. Robert C, Schachter J, Long GV, et al. Pembrolizumab versus ipilimumab in advanced melanoma. *N Engl J Med*. 2015;372:2521-2532.

Together we can reach new heights

Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

Safety and Feasibility of Short-Duration Infusion Obinutuzumab With Venetoclax in Patients With Chronic Lymphocytic Leukemia: A Prospective Observational Study

Presenting Authors: Matthew Lei, PharmD, Uvette Lou, PharmD, Mark Sorial, PharmD, Brenna Rowen, PharmD, E. Bridget Kim, PharmD, Rana Ruchi, PharmD, and Andrea Medrano, BS, PharmD Candidate, Department of Pharmacy, Massachusetts General Hospital, Boston, MA; Priyanka Pullarkat, MD, J. Erika Haydu, MD, PhD, Jeffrey Barnes, MD, Ephraim Hochberg, MD, P. Connor Johnson, MD, Ronald Takvorian, MD, Jeremy Abramson, MD, MS, Jacob D. Soumerai, MD, Julia Lynch, APRN, Emily Patterson, APRN, Katherine Thorp, APRN, Maryanne Sherbourne, APRN, Brianne McGree, APRN, Michelle Knowles, APRN, Josie Ford, BS, Ronald Nemec, BA, Catherine Cho, BA, Center for Lymphoma, Massachusetts General Hospital Cancer Center, Boston, MA; Robert Redd, MS, Department of Data Science, Dana-Farber Cancer Institute, Boston, MA

BACKGROUND: Obinutuzumab is approved in combination with venetoclax for chronic lymphocytic leukemia (CLL)/small lymphocytic lymphoma (SLL) and with chemotherapy for classic follicular lymphoma (cFL). Infusion-related reactions (IRRs) occur in 60% (grade ≥ 3 IRRs: 10%) of patients with CLL/SLL receiving obinutuzumab. During cycles (C) 2 to 6 (C2-6), obinutuzumab was infused over ≥ 195 minutes. A 90-minute short-duration infusion (SDI) obinutuzumab starting with C2 is safe in patients with cFL without grade ≥ 3 IRR in C1. Data on SDI obinutuzumab in CLL/SLL are limited. We report a prospective observational study to evaluate the safety of SDI obinutuzumab in patients with CLL/SLL receiving treatment with venetoclax.

OBJECTIVES: To evaluate the proportion of patients with grade ≥ 3 IRR with 90-minute SDI obinutuzumab in patients with CLL/SLL receiving venetoclax.

METHODS: This prospective, exact single-stage, observational study included the first 28 consecutive patients with treatment naïve or relapsed/refractory CLL/SLL and who received venetoclax identified via a lymphoid malignancy patient registry approved by the institutional review board. Dosing, schedule, premedications, and infusion rates were according to the

obinutuzumab prescribing information. Starting with C2, absent grade ≥ 3 IRR in C1 and an absolute lymphocyte count of $< 5,000/\mu\text{L}$ on day of treatment, obinutuzumab was administered at 100 mg per hour for 30 minutes, then 900 mg per hour for approximately 60 minutes. The primary end point included the proportion of patients with grade ≥ 3 IRR defined per Common Terminology Criteria for Adverse Events v5.0. Response was evaluated per International Workshop on Chronic Lymphocytic Leukemia (iwCLL) 2018.

RESULTS: Between January 23, 2023, and April 11, 2024, 28 consecutive patients with CLL/SLL received venetoclax with SDI obinutuzumab. The median age was 65 years (range, 27-88), and 79% were treatment naïve. Absolute lymphocyte count was $\geq 25,000/\mu\text{L}$ before C1 day (D) 1 in 54% and $\geq 5,000/\mu\text{L}$ before C2D1 in 0%. All patients received SDI obinutuzumab on C2D1. None discontinued obinutuzumab prematurely; the median number of obinutuzumab Cs was 6 (interquartile range, 6-6) and all completed ≥ 4 Cs. All patients achieved an iwCLL objective response. No patients had a grade ≥ 3 IRR. The incidence of grade 1 to 2 IRRs was 53.6%. All IRRs were grade 2 and occurred on C1D1, including 1 patient with recurrent grade 2 IRR on day 2, and none occurred on days 8/15 of C1. One had an IRR after C1; this was a grade 2 IRR occurring on C3D1.

CONCLUSION: SDI obinutuzumab with venetoclax was safe in patients with CLL/SLL. This study met the primary end point and supports adoption of SDI obinutuzumab starting in C2 absent grade ≥ 3 IRR during C1 when absolute lymphocyte count $< 5,000/\mu\text{L}$.

1. Al-Sawaf O, Zhang C, Tandon M, et al. Venetoclax plus obinutuzumab versus chlorambucil plus obinutuzumab for previously untreated chronic lymphocytic leukaemia (CLL14): follow-up results from a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol.* 2020;21: 1188-1200.
2. Al-Sawaf O, Zhang C, Jin HY, et al. Transcriptomic profiles and 5-year results from the randomized CLL14 study of venetoclax plus obinutuzumab versus chlorambucil plus obinutuzumab in chronic lymphocytic leukemia. *Nat Commun.* 2023;14:2147. Erratum in: *Nat Commun.* 2023;14:6724.

3. Townsend W, Hiddemann W, Buske C, et al. Obinutuzumab versus rituximab immunochemotherapy in previously untreated iNHL: final results from the GALLIUM study. *Hemasphere*. 2023;7: e919.
4. Marcus R, Davies A, Ando K, et al. Obinutuzumab for the first-line treatment of follicular lymphoma. *N Engl J Med*. 2017;377:1331-1344.
5. Gazyva (obinutuzumab) injection, for intravenous use [prescribing information] Genentech, Inc.; July 2022. Accessed 10 Oct 2024. (**AU: Access date?**).
www.accessdata.fda.gov/drugsatfda_docs/label/2022/125486s034lbl.pdf
6. Canales MA, Buchholz TA, Bortolini JAP, et al. Obinutuzumab can be administered as a 90-minute short duration infusion in patients with previously untreated follicular lymphoma: GAZELLE end of induction analysis. *Hemasphere*. 2023;7:e860.
7. Fleck C, Karabinos A, Cook A, et al. Evaluation of safety outcomes with transitioning obinutuzumab from standard rate to short duration infusion in patients with chronic lymphocytic leukemia. *Leuk Lymphoma*. 2024;65:1198-1199.
8. Fischer K, Al-Sawaf O, Bahlo J, et al. Venetoclax and obinutuzumab in patients with CLL and coexisting conditions. *N Engl J Med*. 2019;380:2225-2236.

Together we can reach new heights

Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

Updated Biodistribution and Shedding Analysis of RP1 Oncolytic Immunotherapy From the IGNYTE Clinical Trial: Administration and Handling Implications for Oncology Pharmacists

Presenting Authors: Christina Davis, PharmD, University of Colorado Hospital, Aurora, CO; Caroline Robert, MD, PhD, Gustave Roussy and Paris-Saclay University, Villejuif, France; Michael K Wong, MD, PhD, Roswell Park Comprehensive Cancer Center, Buffalo, NY; Mark R Middleton, MD, PhD, FRCP, Churchill Hospital and University of Oxford, Oxford, UK; Mohammed M Milhem, MBBS, Holden Comprehensive Cancer Center, University of Iowa, Iowa City, IA; Joseph J Sacco, PhD, FRCP, The Clatterbridge Cancer Centre, Wirral, UK and University of Liverpool, Liverpool, UK; Judith Michels, MD, PhD, Département de Médecine Oncologique, Gustave Roussy; Gino K In, MD, University of Southern California Norris Comprehensive Cancer Center, Los Angeles, CA; Eva Muñoz Couselo, MD, PhD, Vall d'Hebron Institute of Oncology (VHIO) and Vall d'Hebron Hospital Medical Oncology Department, Barcelona, Spain; Dirk Schadendorf, MD, West German Cancer Center, University Hospital Essen, Essen, Germany; Georgia M Beasley, MD, MHSc, Duke Cancer Institute, Duke University, Durham, NC; Jiaxin Niu, MD, PhD, Banner MD Anderson Cancer Center, Gilbert, AZ; Bartosz Chmielowski, MD, PhD, Jonsson Comprehensive Cancer Center, University of California Los Angeles, Los Angeles, CA; Trisha M Wise-Draper, MD, PhD, University of Cincinnati Cancer Center, University of Cincinnati, Cincinnati, OH; Tawnya Lynn Bowles, MD, Intermountain Medical Center, Murray, UT; Katy K Tsai, MD, Helen Diller Family Comprehensive Cancer Center, University of California San Francisco, San Francisco, CA; Céleste Lebbé, MD, PhD, Université Paris Cité, AP-HP Dermato-Oncology and CIC, Cancer Institute APHP. Nord—Université Paris Cité, INSERM U976, Saint Louis Hospital, Paris, France; Caroline Gaudy-Marqueste, MD, PhD, Aix-Marseille Université, APHM, Centre de Recherche en Cancérologie de Marseille (CRCM), INSERM, U1068, CNRS, UMR7258, UM105, Hôpital Timone, CEPCM, Dermatology and Skin Cancer Department, Marseille, France; Adel Samson, PhD, Leeds Institute of Medical Research at St. James's, University of Leeds, Leeds, UK; Jason A Chesney, MD, PhD, James Graham Brown Cancer Center, University of Louisville, Louisville,

KY; Anna C Pavlick, MD, MBA, Weill Cornell Medical College, New York, NY; Ari M VanderWalde, MD, MPH, FACP, West Cancer Center and Research Institute, Germantown, TN; Kevin Harrington, PhD, The Institute of Cancer Research/Royal Marsden NIHR Biomedical Research Centre, London, UK; Marcus Viana, MD, Laxminarasimha Donthireddy, PhD, Alina Monteagudo, PhD, Junhong Zhu, PhD, Jeannie W Hou, MD, Aaron Clack, PhD, Praveen K Bommareddy, PhD, MS, Replimune, Inc., Woburn, MA

BACKGROUND: RP1 is a herpes simplex virus type 1-based oncolytic immunotherapy currently in clinical development that is administered via intratumoral injection. RP1 plus nivolumab has demonstrated deep and durable responses with a favorable safety profile in advanced melanoma.


OBJECTIVE: To assess the biodistribution and shedding profiles from the skin cancer cohorts of the ongoing IGNYTE clinical trial (NCT03767348), thus informing the administration and handling methodology for RP1.

METHODS: In this open-label, multicenter phase 1/2 study, patients with advanced cancers were treated with the combination RP1 plus nivolumab. After RP1 injections into superficial and deep lesions, injection sites were covered with occlusive dressings. Samples from blood, urine, dressing exteriors, injection sites, oral mucosa, and lesions of suspected herpetic origin were assessed during predose, treatment, and follow-up visits for RP1 DNA by quantitative polymerase chain reaction assay. Positive RP1 DNA swab samples were further assessed by 50% tissue culture infectious dose (TCID₅₀) assay for the presence of live virus.

RESULTS: This analysis included 1573 blood, 1976 urine, 2052 oral mucosa, 1114 dressings, and 1947 injection-site swab samples collected from 278 patients with skin cancer. RP1 DNA was detected in 7.8% of blood, 0.2% of urine, and 18.4% of injection-site swab samples. The incidence of RP1 detection on injection-site dressing exteriors (9.5% of 1114 samples) was lower than that from the injection sites (18.4% of 1947 samples), demonstrating that dressings act as a barrier to RP1 dissemination. RP1 DNA was very rarely present on oral mucosa (0.9% of 2052 samples). At follow-up, RP1 DNA was detected only in injection-site swab samples. All available samples were confirmed to be negative for live virus by TCID₅₀ assay. A total of 8 swab samples from 7 patients from different locations were collected from areas of suspected

herpetic infection; none tested positive for infectious virus. No communal or secondary transmission was reported.

CONCLUSION: RP1 DNA was primarily detected on the immediate surface of injected lesions. Overlying occlusive dressings appear to serve as a protective barrier against RP1 DNA dissemination. No infectious live virus was detected in samples from follow-up visits. These findings indicate that the risk of infection and transmission of RP1 is negligible. Oncolytic immunotherapy such as RP1 is emerging as an integral part of cancer therapeutics. Defining the biodistribution and shedding potential of these agents will be important for pharmacy staff and caregivers to develop the educational activities and best practices necessary to properly handle these agents.



HOPACH

Together we can reach new heights

Completed Research: CLINICAL/TRANSLATIONAL RESEARCH

Venetoclax and Obinutuzumab in Relapsed/Refractory Patients With Chronic Lymphocytic Leukemia or Small Lymphocytic Lymphoma – 3-Year Follow-Up

Presenting Authors: Uvette Lou, PharmD, BCOP, Matthew Lei, PharmD, BCOP, Mark Sorial, PharmD, BCOP, Center for Lymphoma, Massachusetts General Hospital Cancer Center, Boston, MA, and Department of Pharmacy, Massachusetts General Hospital, Boston, MA; Michelle Yu and Andrea Medrano, BS, PharmD Candidate, Center for Lymphoma, Massachusetts General Hospital Cancer Center, Boston, MA, and Northeastern University School of Pharmacy and Pharmaceutical Sciences, Boston, MA; Jeremy S. Abramson, MD, MS, and Jacob D. Soumerai, MD, Center for Lymphoma, Massachusetts General Hospital Cancer Center, Boston, MA [AU: BOSTON CORRECT HERE?]

BACKGROUND: In patients with previously untreated chronic lymphocytic leukemia (CLL) and small lymphocytic leukemia (SLL), venetoclax and obinutuzumab resulted in higher complete response rates compared with chemoimmunotherapy^{1,2} and is FDA approved as first-line therapy but not in relapsed/refractory (R/R) CLL/SLL, where venetoclax plus rituximab is approved despite studies indicating that obinutuzumab is the more effective monoclonal antibody in CLL.^{1,2} Venetoclax and obinutuzumab incorporates an obinutuzumab lead-in phase, which allows for cytoreduction and reduces venetoclax-related tumor lysis syndrome (TLS) risk; whereas in venetoclax plus rituximab, venetoclax starts first, not allowing for cytoreduction. We report 3-year follow-up of our experience with safety and efficacy of venetoclax and obinutuzumab in R/R CLL/SLL.

METHODS: This retrospective study included patients aged ≥ 18 years with R/R CLL who received venetoclax and obinutuzumab at Massachusetts General Hospital locations between July 2019 and June 2022. Updated progression and survival data were collected through October 9, 2024. Safety end points included laboratory or clinical TLS per Howard criteria, infusion-related reactions (IRRs), and hematologic adverse events defined per Common Terminology Criteria for Adverse Events v5.0. Efficacy end points included progression-free survival (PFS) and overall survival (OS), which were estimated using Kaplan-Meier method.

RESULTS: We identified 40 consecutive R/R CLL patients treated with venetoclax and obinutuzumab. Median follow-up was 41 months (range, 2-64). Median age was 72 (range, 51-94) years, with 31 (78%) patients aged ≥ 65 years, 28% (n=11/39) with del(17p)/*TP53* alterations, and 66% (n=21/32) with unmutated *IGHV*. Median previous treatments was 1 (range, 1-6), with 55% (n=22) previously treated with **Bruton tyrosine kinase** inhibitor. Median creatinine clearance (CrCl) was 57 mL/min (range, 22-134), with CrCl < 60 mL/min in 53% (n=21) of patients. Laboratory TLS occurred in 3% (n=1) of patients, arising during obinutuzumab initiation; no patients had clinical TLS. IRRs occurred in 30% (n=12), with 27% (n=11) having grades 1-2 and 3% (n=1) grade 3. Four of 11 (36%) patients who had grade 1-2 IRRs were treated as inpatients. Grade 3/4 hematologic adverse events occurred in 60% (n=24) (neutropenia, 50%; anemia, 25%; thrombocytopenia, 45%). The 3-year PFS and OS rates were 65% (95% confidence interval [CI], 0.51-0.82) and 84% (95% CI, 0.73-0.97), respectively.

CONCLUSION: In 3-year follow-up, venetoclax and obinutuzumab continued to be safe and effective treatment for R/R CLL. Laboratory TLS occurred more frequently during obinutuzumab initiation (3%) than with venetoclax (0%), which may reflect obinutuzumab use over rituximab, or venetoclax and obinutuzumab use in real-world patients with comorbidities that may have excluded them from prospective trials. More intensive TLS mitigation and monitoring should be considered when starting the obinutuzumab phase of venetoclax and obinutuzumab in R/R CLL, including inpatient administration in patients with comorbidities or high tumor burden.

1. Fischer K, Al-Sawaf O, Bahlo J, et al. Venetoclax and obinutuzumab in patients with CLL and coexisting conditions. *N Engl J Med*. 2019;380:2225-2236.
2. Eichhorst B, Niemann C, Kater AP, et al. A randomized phase III study of venetoclax-based time-limited combination treatments (R_{Ve}, G_{Ve}, G_{IVe}) vs standard chemoimmunotherapy (CIT: FCR/BR) in frontline chronic lymphocytic leukemia (CLL) of fit patients: first co-primary endpoint analysis of the international intergroup GAIA (CLL13) trial. *Blood*. 2021;138(suppl 1):71.